

Silver Grove School

Class-12-A

Mathematics

Time Allowed: 3 hours

Maximum Marks: 80

General Instructions:

This Question Paper consists of three sections A, B and C.

*Candidates are required to attempt all questions from **Section A** and all questions*

Section A:** Internal choice has been provided in **two questions of two marks each, two questions of four marks each and two questions of six marks each.

Section B:** Internal choice has been provided in **one question of two marks and one question of four marks.

SECTION A - 65 MARKS

1. In subparts (i) to (x) choose the correct options and in subparts (xi) to (xv), answer the questions as instructed. [15]

- (a) The matrix $\begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$ is [1]

a) a symmetric matrix

b) a unit matrix

c) a diagonal matrix

d) a skew-symmetric matrix

- (b) $\int_a^b \frac{\log x}{x} dx$ is equal to [1]

a) $\frac{\log(b-a)}{b-a}$

b) $\log(a+b) \cdot \log(b-a)$

c) $\log(ab) \cdot \log\left(\frac{b}{a}\right)$

d) $\frac{1}{2} \log(ab) \cdot \log\left(\frac{b}{a}\right)$

- (c) The principal value of $\sin^{-1} \frac{1}{2}$ is [1]

a) Both $\frac{\pi}{6}$ and $\frac{5\pi}{6}$

b) $\frac{\pi}{6}$

c) $\frac{5\pi}{6}$

d) $\frac{-\pi}{6}$

- (d) What is the product of the order and degree of the differential equation $\frac{d^2 y}{dx^2} \sin y + \left(\frac{dy}{dx}\right)^3 \cos y = \sqrt{y}$? [1]

a) 6

b) not defined

- c) 2 d) 3
- (e) A box contains 3 orange balls, 3 green balls and 2 blue balls. Three balls are drawn at random from the box without replacement. The probability of drawing 2 green balls and one blue ball is [1]
- a) $\frac{1}{28}$ b) $\frac{3}{28}$
c) $\frac{167}{168}$ d) $\frac{2}{21}$
- (f) Number of onto (subjective) functions from A to B if $n(A) = 6$ and $n(B) = 3$ are [1]
- a) 340 b) None of these
c) $2^6 - 2$ d) $3^6 - 3$
- (g) If $y = \log\left(\frac{1-x^2}{1+x^2}\right)$ then $\frac{dy}{dx}$ is equal to [1]
- a) $\frac{4x^3}{1-x^4}$ b) $\frac{-4x^3}{1-x^4}$
c) $\frac{1}{4-x^4}$ d) $\frac{-4x}{1-x^4}$
- (h) If $f(x) = |3 - x| + (3 + x)$, where (x) denotes the least integer [1]
- a) neither differentiable nor continuous at $x = 3$ b) continuous but not differentiable at $x = 3$
c) differentiable but not continuous at $x = 3$ d) continuous and differentiable at $x = 3$
- (i) Which of the following is not correct in a given determinant of A, where $A = [a_{ij}]_{3 \times 3}$. [1]
- a) Value of a determinant is obtained by multiplying elements of a row or column by corresponding cofactors
b) Order of minors and cofactors of elements of A is same
c) Order of minor is less than order of the $\det(A)$
d) Minor of an element can never be equal to cofactor of the same element
- (j) **Assertion (A):** If $A = \begin{bmatrix} 2 & 4 \\ 3 & 2 \end{bmatrix}$ and $C = \begin{bmatrix} -2 & 5 \\ 3 & 4 \end{bmatrix}$, then $3A - C = \begin{bmatrix} 8 & 7 \\ 6 & 2 \end{bmatrix}$. [1]
Reason (R): If the matrices A and B are of same order, say $m \times n$, satisfy the commutative law, then $A + B = B + A$.
- a) Both A and R are true and R is the correct explanation of A. b) Both A and R are true but R is not the correct explanation of A.
c) A is true but R is false. d) A is false but R is true.
- (k) If f and g are real function defined by $f(x) = x^2 + 7$ and $g(x) = 3x + 5$, find the $f(t) - f(-2)$ [1]
- (l) Let $A = \begin{bmatrix} 2 & 4 \\ 3 & 2 \end{bmatrix}$, $B = \begin{bmatrix} 1 & 3 \\ -2 & 5 \end{bmatrix}$. Find $A + B$ [1]
- (m) Let g be real function defined by and $g(x) = (x + 4)^3$. Find $\frac{1}{g}$. [1]
- (n) Let E_1 and E_2 be two independent events such that $P(E_1) = p_1$ and $P(E_2) = p_2$. Describe in words of the events whose probability is: $p_1 + p_2 - 2p_1p_2$ [1]
- (o) A bag contains 5 white, 7 red and 8 black balls. Four balls are drawn one by one with replacement, what is the probability that at least one is white? [1]

2. Differentiate $\tan^{-1}\left(\frac{2x}{1-x^2}\right)$ with respect to x , if $x \in (-\infty, -1)$ [2]

OR

Find the points on the curve $y = x^3$ where the slope of the tangent is equal to x -coordinate of the point.

3. Evaluate the Integral: $\int x\sqrt{x^2-1}dx$ [2]

4. Prove that function $f(x) = 3x^5 + 40x^3 + 240x$ is increasing on $\mathbb{R}$. [2]

5. Evaluate: $\int \frac{1}{\sqrt{x}(\sqrt{x}+1)} dx$ [2]

OR

Evaluate: $\int \frac{1}{1+x+x^2+x^3} dx$

6. Let S be the set of all real numbers and let R be a relation in S , defined by $\{R = \{(a, b) : a \leq b^3\}\}$. Show that R satisfies none of reflexivity, symmetry and transitivity. [2]

7. Prove the following. $\cos\left[\sin^{-1}\left(\frac{3}{5}\right) + \cot^{-1}\left(\frac{3}{2}\right)\right] = \frac{6}{5\sqrt{13}}$ [4]

8. Evaluate: $\int \frac{x^2}{(x^4-x^2-12)} dx$. [4]

9. If $y = (\sin^{-1}x)^2$, prove that $(1-x^2) \frac{d^2y}{dx^2} - x \frac{dy}{dx} - 2 = 0$ [4]

OR

If $y = x^n \{a \cos(\log x) + b \sin(\log x)\}$, prove that $x^2 \frac{d^2y}{dx^2} + (1-2n) \frac{dy}{dx} + (1+n^2)y = 0$

10. **Read the text carefully and answer the questions:** [4]

Akash and Prakash appeared for first round of an interview for two vacancies. The probability of Nisha's selection is $\frac{1}{3}$ and that of Ayushi's selection is $\frac{1}{2}$.



- Find the probability that both of them are selected.
- The probability that none of them is selected.
- Find the probability that only one of them is selected.
- Find the probability that atleast one of them is selected.

OR

Read the text carefully and answer the questions:

[4]

A shopkeeper sells three types of flower seeds A_1 , A_2 , A_3 . They are sold in the form of a mixture, where the proportions of these seeds are $4 : 4 : 2$ respectively. The germination rates of the three types of seeds are 45%, 60% and 35% respectively.



Based on the above information:

- Calculate the probability that a randomly chosen seed will germinate.
- Calculate the probability that the seed is of type A2, given that a randomly chosen seed germinates.
- A die is throw and a card is selected at random from a deck of 52 playing cards. Then find the probability of getting an even number on the die and a spade card.
- If A and B are any two events such that $P(A) + P(B) - P(A \text{ and } B) = P(A)$, then find $P(A|B)$.

11. **Read the text carefully and answer the questions:**

[6]

A trust fund has ₹ 35000 that must be invested in two different types of bonds, say X and Y. The first bond pays 10% interest p.a. which will be given to an old age home and second one pays 8% interest p.a. which will be given to WWA (Women Welfare Association).

Let A be a 1×2 matrix and B be a 2×1 matrix, representing the investment and interest rate on each bond respectively.



Based on the above information, answer the following questions.

- If ₹ 15000 is invested in bond X, then what is the matrix representation of A and B?
- If ₹ 15,000 is invested in bond X, how can we determine the total amount of interest received on both bonds?
- How much is the investment in two bonds if the trust fund obtains an annual total interest of ₹3200?

12. a. Solve the differential equation for a particular solution:

[6]

$$dy = (5x - 4y) dx, \text{ when } y = 0 \text{ and } x=0$$

- b. Solve the differential equation $ydx - (x + 2y^2) dy = 0$

OR

Solve the diff. eq $\frac{dy}{dx} + \frac{y^2+y+1}{x^2+x+1} = 0$

13. The sum of the surface areas of a cuboid with sides x , $2x$ and $\frac{x}{3}$ and a sphere is given to be constant. Prove that the sum of their volumes is minimum, if x is equal of three times the radius of sphere. Also, find the minimum value of the sum of their volumes.

[6]

OR


Which of the following functions are decreasing on $(0, \frac{\pi}{2})$?

- $\cos x$
- $\cos 2x$
- $\cos 3x$
- $\tan x$

14. **Read the text carefully and answer the questions:**

[6]

Recent studies suggest that roughly 12% of the world population is left handed.



Depending upon the parents, the chances of having a left handed child are as follows:

A. When both father and mother are left handed:

Chances of left handed child is 24%.

B. When father is right handed and mother is left handed:

Chances of left handed child is 22%.

C. When father is left handed and mother is right handed:

Chances of left handed child is 17%.

D. When both father and mother are right handed:

Chances of left handed child is 9%.

Assuming that $P(A) = P(B) = P(C) = P(D) = \frac{1}{4}$ and L denotes the event that child is left handed.

- Find $P\left(\frac{L}{C}\right)$
- Find $P\left(\frac{L}{A}\right)$
- Find $P\left(\frac{A}{L}\right)$
- Find the probability that a randomly selected child is left handed given that exactly one of the parents is left handed.

SECTION B - 15 MARKS

15. In subparts (i) and (ii) choose the correct options and in subparts (iii) to (v), answer the questions as instructed. [5]

(a) The vectors $3\hat{i} - \hat{j} + 2\hat{k}$, $2\hat{i} + \hat{j} + 3\hat{k}$ and $\hat{i} + \lambda\hat{j} - \hat{k}$ are coplanar if the value of λ is: [1]

- 2
- 0
- 2
- Any real number

(b) If a line makes an angle of 30° , 60° , 90° with the positive direction of x, y, z-axes, respectively, then find its direction cosines. [1]

(c) Find the value of $(\hat{i} \times \hat{j}) \cdot \hat{k} + \hat{i} \cdot \hat{j}$. [1]

(d) Determine whether the given plane $2x - 2y + 4z + 5 = 0$ and $3x - 3y + 6z - 1 = 0$ are parallel or perpendicular, and in case they are neither, find the angles between them. [1]

- The planes are at 45°
- The planes are parallel
- The planes are at 55°
- The planes are perpendicular

(e) Find the direction cosines of the line joining the points P(4, 3, -5) and Q(-2, 1, -8). [1]

16. If $\vec{a}, \vec{b}, \vec{c}$ are three mutually perpendicular unit vectors, then prove that $|\vec{a} + \vec{b} + \vec{c}| = \sqrt{3}$ [2]

OR

If $\vec{a} = (\hat{i} - 2\hat{j})$, $\vec{b} = (2\hat{i} - 3\hat{j})$ and $\vec{c} = (2\hat{i} + 3\hat{k})$, find $(\vec{a} + \vec{b} + \vec{c})$.

17. Write the value of k for which the planes $x - 2y + kz = 4$ and $2x + 5y - z = 9$ are perpendicular. [4]

OR

Find the angle between the lines whose direction cosines are given by the equations: $3l + m + 5n = 0$ and $6mn - 2nl +$

$$5lm = 0.$$

18. Using integration, find the area of the region bounded by the line $y - 1 = x$, the x -axis and the ordinates $x = -2$ and $x = 3$. [4]